A simple proof that π is irrational

Ivan Niven

Let $\pi = a/b$, the quotient of positive integers. We define the polynomials

$$f(x) = \frac{x^n (a - bx)^n}{n!},$$

$$F(x) = f(x) - f^{(2)}(x) + f^{(4)}(x) - \dots + (-1)^n f^{(2n)}(x),$$

the positive integer n being specified later. Since n!f(x) has integral coefficients and terms in x of degree not less than n, f(x) and its derivatives $f^{(j)}(x)$ have integral values for x = 0; also for $x = \pi = a/b$, since f(x) = f(a/b - x). By elementary calculus we have

$$\frac{\mathrm{d}}{\mathrm{d}x}\left\{F'(x)\sin x - F(x)\cos x\right\} = F''(x)\sin x + F(x)\sin x = f(x)\sin x$$

and

(1)
$$\int_0^{\pi} f(x) \sin x \, dx = \left[F'(x) \sin x - F(x) \cos x \right]_0^{\pi} = F(\pi) + F(0).$$

Now $F(\pi) + F(0)$ is an integer, since $f^{(j)}(\pi)$ and $f^{(j)}(0)$ are integers. But for $0 < x < \pi$,

$$0 < f(x)\sin x < \frac{\pi^n a^n}{n!},$$

so that the integral in (1) is positive, but arbitrarily small for n sufficiently large. Thus (1) is false, and so is our assumption that π is rational.

PURDUE UNIVERSITY

Received by the editors November 26, 1946, and, in revised form, December 20, 1946.